



PATIENT PRESENTING CLINICAL SIGNS

Casey O'Brien Left forelimb lameness, collapsing towards the left intermittently.
 Abnormal PE/Chem/CBC/UA Results: Ambulatory x 4 but noticeable weight-bearing lameness
 Left forelimb.

SPECIES COMPUTED TOMOGRAPHY OF THE FRONT LIMBS

Canine A high resolution pre- and post-contrast CT study of the front limbs is provided for review.

COMPUTED TOMOGRAPHIC FINDINGS

BREED Multifocal spondylosis formation is seen along the thoracic spine.

Labrador Retriever The periarticular bones of both shoulder joints present mild osteophyte new bone formation.

SEX

Spayed Female The periarticular bones of the left elbow joint present advanced osteophyte new bone formation. The medial coronoid process of the left elbow joint is irregular marginate. At laterocranial aspect of the medial coronoid process of the left elbow joint, three osseous fragments are measuring up to between 0.6 to 2.0 mm in size. A mild radio-ulnar step formation of 2.5 mm is seen. The left elbow joint presents a moderate thickening of the joint capsule.

AGE

9 Years The periarticular bones of the right elbow joint present mild to moderate osteophyte new bone formation. The medial coronoid process of the right elbow joint presents a heterogeneous density with a hypoattenuating region in the lateral aspect.

COMPUTED TOMOGRAPHIC DIAGNOSIS

INTERPRETED BY

Sebastian Schaub, DVM
 Dr. med. vet. DipECVDI

- Fragmented medial coronoid process (FCP) left elbow joint and incongruity of the left elbow joint
- Advanced degenerative osteoarthritis left elbow joint
- Coronoid disease right elbow joint
- Mild to moderate degenerative osteoarthritis right elbow joint
- Mild degenerative osteoarthritis shoulder joints bilaterally
- Spondylosis deformans

HOSPITAL NAME

Mobile Pet Imaging

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

REFERRING VET

Meaux

The CT study is consistent with advanced degenerative joint disease of the left elbow joint due to FCP, the right elbow joint presents evidence of mild degenerative joint disease due to underlying coronoid disease.

Arthroscopy of the left elbow joint might be considered to revise the elbow joint and remove the fragments. However, the degenerative changes are irreversible, and lameness is likely to persist after surgical intervention, warranting conservative treatment options.

INVOICE

52978

DATE

7-19-22



PATIENT

Casey O'Brien

SPECIES

Canine

BREED

Labrador Retriever

SEX

Spayed Female

AGE

9 Years

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVCI

HOSPITAL NAME

Mobile Pet Imaging

REFERRING VET


Meaux

INVOICE

52978

DATE

7-19-22





PATIENT

Casey O'Brien

SPECIES

Canine

BREED

Labrador Retriever

SEX

Spayed Female

AGE

9 Years

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

HOSPITAL NAME

Mobile Pet Imaging

REFERRING VET

Meaux

The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Sebastian Schaub, Sebastian Schaub, DVM, Dr. med. vet. DipECVDI
sebast.schaub@gmail.com

INVOICE

52978

DATE

7-19-22

