


PATIENT PRESENTING CLINICAL SIGNS

Max Tugwell 3 weeks ago, he was here and said he looks great and then he started limping that night. Max won't walk very far, he limps on his left leg a lot. He will just plop down and not want to go on very long walks. He had surgery a year ago on his leg and he won't use it very much. He'll just walk a couple of feet and just sits down. Been giving pain medication every night, seems to help but he won't walk.

SPECIES

Canine

BREED

Labrador Retriever

SEX

Neutered Male

RADIOGRAPHIC STUDY OF THE LUMBAR SPINE, PELVIS AND STIFLE JOINTS

A complete set of radiographs of the lumbar spine, pelvis and the stifle joints is provided for review.

AGE

7 Years 6 Months

The osseous and surrounding soft tissue structures of the lumbar spine present without abnormalities. The osseous and surrounding soft tissue structures of the pelvis are within normal limits. Both coxofemoral joints present smooth osseous margins and congruent joint spaces.

INTERPRETED BY

Sebastian Schaub,
 DVM Dr. med. vet.
 DipECVDI

The periarticular bones of the left stifle joint present moderate osteophyte new bone formation. A mild intracapsular soft tissue swelling of the left stifle joint is appreciated. A TPLO implant is appreciated at the medioproximal aspect of the left tibia, being unremarkable in place. The osteotomy of the proximal tibia is filled with trabecular bone.

HOSPITAL NAME

Elizabeth AH

The periarticular bones of the right stifle joint are smooth, there is no signs for intracapsular swelling. A subcutaneous fat opaque swelling is seen at the mediodistal aspect of the right stifle joint.

RADIOGRAPHIC DIAGNOSIS
REFERRING VET

Dr. Leon Anderson

- History of surgical management of cranial cruciate ligament pathology of the left stifle joint via TPLO – the osteotomy is in the remodeling phase
- Degenerative osteoarthritis left stifle joint
- Mild articular swelling left stifle joint
- Suspect subcutaneous lipoma medial aspect right stifle joint
- Normal lumbar spine
- Normal pelvis

INVOICE

44592

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS
DATE

8/10/23

An underlying cause for the ongoing left hind limb lameness is not appreciated, but the degenerative joint disease secondary to pathology of the cranial cruciate ligament. A consideration might be meniscal pathology. Removing the TPLO implant might be beneficial as well.



PATIENT

Max Tugwell

SPECIES

Canine

BREED

Labrador Retriever

SEX

Neutered Male

AGE

7 Years 6 Months



INTERPRETED BY

Sebastian Schaub,
DVM Dr. med. vet.
DipECVDI

The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

HOSPITAL NAME

Elizabeth AH

Sebastian Schaub, Sebastian Schaub, DVM, Dr. med. vet. DipECVDI
sebast.schaub@gmail.com

REFERRING VET

Dr. Leon Anderson

INVOICE

44592

DATE

8/10/23