



PATIENT PRESENTING CLINICAL SIGNS

Mao Mao Le possible infiltrative lipoma between the shoulder blades. There were 2 masses initially noticed 2 months ago. The smaller mass on the right side of the shoulder blade was initially 1.5 cm and hasn't grown larger, the one between shoulder blades appears to have grown rapidly to 2cm within 2 months. a fine needle aspirate was performed on one mass (the smaller one) - cytology revealed adipose tissue. Cytology of the other mass not been completed.

SPECIES

Feline

COMPUTED TOMOGRAPHY OF THE THORAX

A high resolution pre- and post-contrast CT study of the thorax in a lung and soft tissue reconstruction is provided for review.

BREED

Domestic Shorthair

COMPUTED TOMOGRAPHIC FINDINGS

The left thyroid gland is prominent.

SEX

Level with the spinous process of T4&T5, the subcutaneous fat at the right dorsal aspect of the thoracic spine presents a mild increased volume, demarcated by a very thin soft tissue attenuating capsule; measuring 27 x 10 x 24 mm in size.

MN

AGE

The sternal, cranial mediastinal and tracheobronchial lymph nodes are small elongated with a normal short-to-long-axis-ratio is < 0.5, the attenuation and contrast enhancement pattern is uniform and considered within normal limits.

8 Years

The cardiovascular structures including the pulmonary vasculature are within normal limits.

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

The bronchial tree presents with regular branching and tapers uniformly towards the periphery as expected, the bronchial walls are thin and smooth. The bronchus-to-artery ratio is within normal limits.

The volume of the lung is mildly decreased, and multiple ground glass attenuating regions of the lung parenchyma are appreciated.

HOSPITAL NAME

Animal Health
Partners

Small incidental gas pockets are seen within the esophageal lumen, there is no evidence of abnormal dilation.

COMPUTED TOMOGRAPHIC DIAGNOSIS

REFERRING VET

Dr. Debbie Reynolds

- Subcutaneous lipoma right dorsal aspect thoracic spine
- Regions of dystelectasis of the lung parenchyma
- Nodular enlargement left thyroid gland
- No evidence of pulmonary metastatic disease

INVOICE

53744


INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

There is a lipoma at the right dorsal aspect of the thoracic spine, level with T4/T5, without overt signs of local infiltrative growth. Complete surgical excision is considered feasible.

DATE

8-25-22

The nodular enlargement of the left thyroid gland can present (non)functional nodular hyperplasia or adenoma formation. Correlate with T4 values.



PATIENT

Mao Mao Le

SPECIES

Feline

BREED

Domestic Shorthair

SEX

MN

AGE

8 Years

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

HOSPITAL NAME

Animal Health
Partners

REFERRING VET


Dr. Debbie Reynolds


INVOICE

53744

DATE

8-25-22





PATIENT

Mao Mao Le

SPECIES

Feline

The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

BREED

Domestic Shorthair

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Sebastian Schaub, Sebastian Schaub, DVM, Dr. med. vet. DipECVDI
sebast.schaub@gmail.com

SEX

MN

AGE

8 Years

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

HOSPITAL NAME

Animal Health
Partners

REFERRING VET

Dr. Debbie Reynolds

INVOICE

53744

DATE

8-25-22