



PATIENT

Ruger Ross

PRESENTING CLINICAL SIGNS

Left forelimb chronic lameness. Holds leg out straight when running. Reluctant to bend at the elbow. No pain noted on flexion of carpus, elbow or scapulohumeral joints. Radiographs of both the left and the right forelimb are unremarkable.

SPECIES

Canine

Abnormal PE/Chem/CBC/UA Results: wnl

COMPUTED TOMOGRAPHY OF THE FRONT LIMBS

A high resolution pre- and post-contrast CT study of the front limbs is provided for review.

BREED

Golden Retriever

COMPUTED TOMOGRAPHIC FINDINGS

Parts of the right shoulder joint are cropped by the field of view.

Both shoulder joints present smooth osseous margins and the surrounding soft tissue structures are within normal limits.

SEX

Male Intact

The medial coronoid process of the left elbow joint presents mild undulating margins. The tip of the medial coronoid process of the left elbow joint has a heterogeneous density with generalized decreased attenuation behavior. At the caudal aspect of the cranial tip of the medial coronoid process of the left elbow joint, an incomplete thin fissure transverse fissure line is appreciated.

AGE

2

Caudal to the cranial tip of the medial coronoid process of the right elbow joint, a mild oblique fissure line is demarcating an osseous fragment, measuring 3.5 x 3.5 x 4.0 mm in size.

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

COMPUTED TOMOGRAPHIC DIAGNOSIS

- Coronoid disease elbow joints bilaterally
- Structural normal shoulder joints

HOSPITAL NAME

Advanced Animal
Imaging

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

The findings are consistent with pathology of the medial coronoid process of both elbow joints – suspect non-vital tip with fissure lines. At this point there is no evidence of accompanying degenerative joint disease.

REFERRING VET

Blair Hollowell


Arthroscopy of both or at least the left elbow joint is recommended to revise the elbow joints and remove the potential fragments of the medial coronoid process. Overall prognosis for the patient to be free of lameness is considered good after fragment removal. However, episodes of lameness may reoccur after exercise.

INVOICE

53792

DATE

8-29-22



PATIENT

Ruger Ross

SPECIES

Canine

BREED

Golden Retriever

SEX

Male Intact

AGE

2

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

HOSPITAL NAME

Advanced Animal
Imaging

REFERRING VET



Blair Hollowell


INVOICE

53792

DATE

8-29-22





PATIENT

Ruger Ross

The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

SPECIES

Canine

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

BREED

Golden Retriever

Sebastian Schaub, Sebastian Schaub, DVM, Dr. med. vet. DipECVDI
sebast.schaub@gmail.com

SEX

Male Intact

AGE

2

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

HOSPITAL NAME

Advanced Animal
Imaging

REFERRING VET

Blair Hollowell

INVOICE

53792

DATE

8-29-22