



PATIENT PRESENTING CLINICAL SIGNS

Teddy Young presented for TPLO - elevated alkp - abdo scan: gastric nodule noted
Abnormal PE/Chem/CBC/UA Results: alkp/gluc elevated

SPECIES COMPUTED TOMOGRAPHY OF THE ABDOMEN

Canine A pre- and post-contrast CT study of the abdomen in a bone and soft tissue reconstruction is provided for review.

BREED COMPUTED TOMOGRAPHIC FINDINGS

Cairn Terrier The serosal fat presents normal attenuation behavior. There is no evidence of peritoneal effusion or peritonitis.

Both kidneys present within normal limits for size, shape and organ architecture. After contrast administration a bilaterally symmetric and uniform nephro- and pyelogram is noted.

SEX The adrenal glands are within normal limits for size, shape and organ architecture.

MN Both liver and spleen present with normal shape, even surface, uniformly attenuating parenchyma and homogeneous contrast enhancement, unremarkable.

AGE The portal vein presents a normal order of its tributary veins and intrahepatic branching. No abnormal vessel is noted inside and outside of the liver parenchyma.

8 (approx.) The pancreas is evenly contoured, the pancreatic parenchyma is homogeneous and presents uniform contrast enhancement.

INTERPRETED BY

Sebastian Schaub, DVM Dr. med. vet. DipECVDI Level with the lesser curvature of the stomach, a mild hypoattenuating intramural mass is seen, presenting mild punctuate mineralization; measuring 12 x 12 x 15 mm in size. The wall layering of the gastric wall appears to be maintained, level with the intramural mass.

HOSPITAL NAME The bony and surrounding soft tissue structures reveal no abnormalities.

Advanced Veterinary Imaging **COMPUTED TOMOGRAPHIC DIAGNOSIS**

- Intramural gastric mass, lesser curvature

REFERRING VET INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

Eamon The CT study is supporting the diagnosis of an intramural mass of the lesser curvature with leiomyoma/-sarcoma being on the top of the list, other differentials include adenocarcinoma, round cell tumor, other. Surgical excision of the mass is considered feasible.

INVOICE The CT study of the abdomen presents no macromorphological abnormalities, explaining the elevated liver enzymes. Ultrasound guided FNA sampling and/or Tru-cut biopsy can be used as minimally invasive methods for further workup.

DATE

8-4-22



PATIENT

Teddy Young

SPECIES

Canine

BREED

Cairn Terrier

SEX

MN

AGE

8 (approx.)

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

HOSPITAL NAME

Advanced Veterinary
Imaging

REFERRING VET

Eamon

INVOICE

53286

DATE

8-4-22



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Sebastian Schaub, Sebastian Schaub, DVM, Dr. med. vet. DipECVDI
sebast.schaub@gmail.com