



PATIENT PRESENTING CLINICAL SIGNS

Mimi Airgood History: this patient has audible breathing, most likely from upper airway. R/o pneumonia or other
Abnormal PE/Chem/CBC/UA Results: n/a

SPECIES COMPUTED TOMOGRAPHIC STUDY OF THE THORAX

Canine A pre- and post-contrast CT study of the thorax in a bone and soft tissue reconstruction is provided for review.

BREED COMPUTED TOMOGRAPHIC FINDINGS

Chihuahua Mild to moderate motion artefacts are present.

SEX

Female

The bony and surrounding soft tissue structures are within normal limits.

The mediastinum contains a moderate amount of fat. The sternal, cranial mediastinal and tracheobronchial lymph nodes are small elongated with a normal short-to-long-axis-ratio is < 0.5, the attenuation and contrast enhancement pattern is uniform and considered within normal limits.

AGE

12 Years

The cardiovascular structures including the pulmonary vasculature are within normal limits.

The bronchial tree presents with regular branching and tapers uniformly towards the periphery as expected, the bronchial walls are thin and smooth. The bronchus-to-artery ratio is within normal limits.

INTERPRETED BY

Sebastian Schaub,
DVM Dr. med. vet.
DipECVDI

The lung parenchyma presents the expected architecture and attenuation behavior.

Small incidental gas pockets are seen within the esophageal lumen, there is no evidence of abnormal dilation.

HOSPITAL NAME

Green Dog Dental
and Wellness

Throughout the hepatic parenchyma, multiple mild hypoattenuating parenchymal lesions are appreciated.

COMPUTED TOMOGRAPHIC DIAGNOSIS

REFERRING VET

Dr. Lena Horn

- Pre- and post-contrast hypoattenuating hepatic parenchymal lesions
- Normal thorax

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

The CT study of the thorax presents without pathology of the lower airways.

INVOICE

16759

The most likely differentials for the hepatic parenchymal hypoattenuating lesions are hepatic cysts, nodular hyperplasia/regeneration nodules, steroid induced hepatopathy, hepatitis or less likely neoplastic infiltration. Ultrasound guided FNA sampling can be used as advanced minimally invasive diagnostic tool.

DATE

8/5/22



PATIENT

Mimi Airgood

SPECIES


Canine

BREED

Chihuahua

SEX

Female



AGE

12 Years

The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

INTERPRETED BY

Sebastian Schaub,
DVM Dr. med. vet.
DipECVDI

Sebastian Schaub, Sebastian Schaub, DVM, Dr. med. vet. DipECVDI
sebast.schaub@gmail.com

HOSPITAL NAME

Green Dog Dental
and Wellness

REFERRING VET

Dr. Lena Horn

INVOICE

16759

DATE

8/5/22