



PATIENT PRESENTING CLINICAL SIGNS

Bone Inman
 Not defecating for 4 days, pooped this morning. Urinating less for past few days.
 Abnormal PE/Chem/CBC/UA Results: Hydration: Appropriately hydrated Mentation: BAR, swatting and attacking prior to being anesthetized EENT: No nasal discharge; clear no discharge OU; clean no debris AU; No cough on tracheal palpation. Oral Cavity: No dental tartar present
SPECIES Lymph Nodes: Symmetrical, no changes in size, shape, consistency Skin: Good hair coat, no signs of ectoparasites. No lesions noted. CV/Respiratory: No murmur or crackles/wheezing auscultated. Synchronous pulses, normal rate. Normal bronchovesicular sounds. Abd/GI: Soft non painful abdomen, no organomegaly, no abnormalities on abdominal palpation Uro/Perineum: N
 Feline Musculoskeletal: Ambulatory x4, no lameness noted. No pain on palpation of limbs. BCS 5/9
BREED Neurological: Appropriate
 DSH

RADIOGRAPHIC STUDY OF THE ABDOMEN

Radiographs of the abdomen in two imaging planes are provided for review.

SEX RADIOGRAPHIC FINDINGS

MN
 The surrounding bony structures are within normal limits.

AGE
 No abnormalities of the extraabdominal soft tissues are noted. The abdominal wall is smooth and thin.

6 Years
 Two eggshell like mineralized bodies are superimposed on the right & left mid ventral abdomen; measuring 15 mm in size.

INTERPRETED BY

Sebastian Schaub, DVM
 Dr. med. vet. DipECVDI

The serosal fat medial to the spleen presents a mild soft tissue striation.

The liver is appropriate in position, size and presents uniform opacity.

The splenic head is in the anticipated position and within normal limits for size and opacity.

HOSPITAL NAME

DPC Veterinary
 Hospital

Both kidneys are seen and present with normal size, shape, delineation and opacity. The urinary bladder is in its anticipated position. No radiopaque calculi are noted throughout the upper and lower urinary tract.

The stomach is in its anticipated position and presents normal content.

REFERRING VET

White

The small intestinal loops are of even diameter and non-dilated, a small amount of gas is seen within the small intestinal loops and considered within normal limits.

The colon is seen in the expected position and presents with appropriate content.

RADIOGRAPHIC DIAGNOSIS

- INVOICE**
- Focal soft tissue striation peritoneal fat
 - Two Bates bodies (nodular fat necrosis) in the ventral abdomen, incidental
- 54003

DATE

9-7-22

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

The focal soft tissue striation of the peritoneal fat medial to the spleen can be caused by the left lobe of the pancreas or presents focal peritonitis – e.g. secondary to pancreatitis. If not done so yet recommend complementing workup by complete blood work including fplli. An abdominal



PATIENT

ultrasound examination can be used as advanced imaging modality as well.

Bone Inman

SPECIES

Feline

BREED

DSH

SEX

MN

AGE

6 Years

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

HOSPITAL NAME

DPC Veterinary
Hospital



REFERRING VET

White

The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

INVOICE

54003

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Sebastian Schaub, Sebastian Schaub, DVM, Dr. med. vet. DipECVDI
sebast.schaub@gmail.com

DATE

9-7-22