



PATIENT PRESENTING CLINICAL SIGNS

Gizmo Barrs Gizmo presented for acute lethargy, inappetance 1 day duration, 1 episode of vomiting. Xrays were taken to evaluate further in consideration of possible gastric FB and explore
Abnormal PE/Chem/CBC/UA Results: Labs were wnl

SPECIES RADIOGRAPHIC STUDY OF ABDOMEN

K9 Orthogonal views of the abdomen are available for review totaling 3 images. Two lateral views, right and left. One ventrodorsal view.

BREED RADIOGRAPHIC FINDINGS

Shih Tzu The stomach is variably distended. There is a moderate amount of gas admixed with homogeneous and heterogeneous soft tissue opaque material; however, the soft issue material is mobile, and it is not visibly retained in the pyloric antrum.

SEX
M The duodenum is not clearly defined, and it is possible due to soft tissue intraluminal fluid retention and/or mild loss of adjacent serosal detail.

AGE
1 There is regionally reduced abdominal serosal detail in the right cranial abdominal region, at the right 11th and 13th intercostal spaces, border effacing the pylorus, proximal duodenum, and transverse colon. The remaining abdominal serosal detail is normal.

The small intestines are normally distributed, of normal diameter, contain a moderate amount of gas and homogeneous soft tissue fluid material. No radiopaque foreign material is noted.

INTERPRETED BY
The colon contains a moderate amount of heterogeneous fecal material and gas.

Tilde Rodrigues Froes, DMV, MSc., Dr. Med Vet., Dipl. CBraRVet The liver, renal silhouettes, and spleen are normal in size, shape, and contour.

The urinary bladder is mildly filled with the apex in the plane of cranial L5 and is homogenously soft tissue opaque.

HOSPITAL NAME
The collimated musculoskeletal structures and thorax are normal.

Golden Isles Animal Hospital

RADIOGRAPHIC DIAGNOSIS

- Gastric ileus with moderate gas, fluid, and post-prandial content. Differential diagnosis, less likely partial obstruction, or intermittent pyloric outflow obstruction.
- Mildly reduced abdominal serosal detail in the right cranial quadrant. The differential diagnosis includes regional peritonitis, nonobstructive gastroenteritis, and colitis correlated to pancreatitis.
- Otherwise, normal abdomen.

INVOICE INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

59090 The stomach is moderately distended, and the duodenum is not clearly defined. This could be correlated to gastric ileus. However, a partial or intermittent outflow obstruction cannot be completely excluded but is less likely. If the patient has persistent vomiting, an abdominal

DATE

6-29-23



PATIENT ultrasonography, as mentioned, is suggested for better evaluation of the antrum, pyloric region, duodenum, and pancreas. Another differential diagnosis includes an unspecific gastroenteritis.

Gizmo Barrs

SPECIES

K9

BREED

Shih Tzu

SEX

M

AGE

1

INTERPRETED BY

Tilde Rodrigues Froes,
DMV, MSc., Dr. Med
Vet., Dipl. CBraRVet

HOSPITAL NAME

Golden Isles Animal
Hospital

REFERRING VET



James Hornbuckle

INVOICE

59090

DATE

6-29-23





PATIENT

Gizmo Barrs

SPECIES

K9

BREED

Shih Tzu

SEX

M

AGE

1

INTERPRETED BY

Tilde Rodrigues Froes,
DMV, MSc., Dr. Med
Vet., Dipl. CBraRVet

Tilde Rodrigues Froes, DMV, MSc., Dr. Med.Vet., Dipl.CBraRVet
info@sonopath.com

HOSPITAL NAME

Golden Isles Animal
Hospital

REFERRING VET


James Hornbuckle

INVOICE

59090

DATE

6-29-23



The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.