

**PATIENT**

Gracie McCleary

**SPECIES**

Feline

**BREED**

DSH

**SEX**

Female Spayed

**AGE**

7.5 years

**WEIGHT**

11.3 lbs

**INTERPRETED BY**

Tam Mengine, DVM,  
DABVP (canine/feline  
practice)

**IMAGING  
PERFORMED BY**

Dr. Tam Mengine

**HOSPITAL NAME**

Stoney Creek VH

**REFERRING VET**

Dr. Tam Mengine

**INVOICE**

12993

**DATE**

5.9.23

**PRESENTING CLINICAL SIGNS**

History: Two-year history of intermittent diarrhea and weight loss, also very itchy ears and sneezing. CBC / Chem / U/A / T4 and FPL have been normal.

**ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN**

**Urinary System**

The urinary bladder is moderately distended with anechoic urine, and no luminal sediment is present. The ureteral papillae, trigone and pelvic urethra are of normal appearance, and the ureters are not visible (normal). No masses, calculi or mucosal irregularities are noted. Urethra visualized to 3.0 cm.

Both kidneys exhibit mildly decreased corticomedullary differentiation. Infarcts are seen within the renal cortex of the left kidney. There is no evidence of nephrolithiasis, pyelectasia or hydronephrosis. The proximal ureters are not visible (normal). The left kidney is 3.3 cm in length. The right kidney is 3.7 cm in length.

**Adrenal Glands**

The adrenal glands are both identified in their normal locations. They are normal in size and shape with appropriate parenchymal echogenicity and normal phrenic vasculature. The left adrenal gland height is 3.6 mm at the caudal pole. The right adrenal gland height 3.8 mm at the caudal pole.

**Spleen**

The spleen is of appropriate size and has a normal, homogenous parenchyma with a smooth, continuous capsular surface. The splenic vasculature is normal with no evidence of congestion or thrombosis, and blood flow through the splenic hilus appears normal. Thickness at the splenic hilus is normal at 9.2 mm.

**Liver**

The liver is of appropriate size and shape, with sharp borders and a mildly coarse parenchymal echotexture that is hypoechoic to the spleen. The portal and hepatic vasculature are of normal size and appearance with no evidence of congestion or thrombosis.

The gallbladder is moderately distended with anechoic contents. The wall was thin and continuous with no focal lesions. The cystic and common bile ducts are normal / not visible.

**Gastrointestinal**


The stomach is empty. The gastric wall is subjectively normal in thickness, and exhibits appropriate wall layering, but cannot be accurately measured due to normal deviations of the rugal folds. The pylorus is of normal appearance.

The small bowel has diffuse changes to the normal 1:3 muscularis to mucosa ratio. Wall measurements are increased up to 2.4 mm for duodenum and 2.7 mm for jejunum. Overall wall layering is preserved. Intestinal motility appears normal.

The visible portions of the colon are of normal thickness, up to 1.0 mm, with intact wall layering. The ileocecal junction is visualized and appears normal.

**Pancreas**

The areas of the limbs and body of the pancreas are isoechoic to the surrounding mesenteric fat, with normal capsular appearance. There is no evidence of peripancreatic inflammation. The pancreatic duct appears normal.



**PATIENT**

Gracie McCleary

**Free Abdomen**

There is no evidence of free fluid within the peritoneal cavity. The omentum and intra-abdominal fat are of appropriate echogenicity. Enlarged abdominal lymph nodes are not observed. The aortic trifurcation has normal blood flow with no evidence of thrombosis.

**SPECIES**

Feline

**ULTRASONOGRAPHIC FINDINGS**

**Primary Findings**

- Diffuse small bowel changes, typical of infiltrative bowel disease

**BREED**

DSH

**Secondary Findings**

- Mild chronic renal changes with an infarct in the left kidney

**SEX**

Female Spayed

**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

The changes in the gastrointestinal tract are suggestive of infiltrative bowel disease, including both inflammatory bowel disease or low grade gastrointestinal lymphoma. Recommendations include:

**AGE**

7.5 years

- Fecal parasite testing and empiric fenbendazole treatment
- Trials with a novel protein or hydrolyzed diet
- A complete GI panel, or empiric cobalamin supplementation
- Empiric therapy with prednisolone at 2-4mg / kg daily could be considered if a diet trial is unsuccessful.
- Definitive diagnosis would require biopsy of the affected tissue, ideally with intra-operative ultrasonographic guidance. If there is concurrent lymphadenopathy, ultrasound-guided sampling of the lymph node using a 25 or 22G needle could be considered.

**WEIGHT**

11.3 lbs

**INTERPRETED BY**

Tam Mengine, DVM,  
DABVP (canine/feline  
practice)

The infarct in the kidney is likely an incidental finding, however, if signs of cardiac disease are noted (such as a murmur or other symptoms) then an echocardiogram would be recommended to rule out an underlying cardiomyopathy.

**IMAGING PERFORMED BY**

Dr. Tam Mengine



**HOSPITAL NAME**

Stoney Creek VH

**REFERRING VET**

Dr. Tam Mengine



**INVOICE**

12993

**DATE**

5.9.23



**PATIENT**

Gracie McCleary

**SPECIES**

Feline

**BREED**

DSH

**SEX**

Female Spayed

**AGE**

7.5 years

**WEIGHT**

11.3 lbs

**INTERPRETED BY**

Tam Mengine, DVM,  
DABVP (canine/feline  
practice)

**IMAGING  
PERFORMED BY**

Dr. Tam Mengine

**HOSPITAL NAME**

Stoney Creek VH

**REFERRING VET**


Dr. Tam Mengine

**INVOICE**

12993

**DATE**

5.9.23



The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance, please contact me.

**Tam Mengine, DVM, DABVP (canine/feline practice) info@SonoPath.com**