

**PATIENT PRESENTING CLINICAL SIGNS**

Savy Busch P has been vomiting and has had diarrhea since July 19 - blood present in both on and off. P lethargic. Has been eating chicken and rice. Most recent exam (July 28): BAR but euhydrated dog with blood in stool like gel.

**SPECIES**

Canine

Abnormal PE/Chem/CBC/UA Results: Negative diarrhea panel. Lab work wnl.

**BREED**

Weimaraner X

**ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN**

**Urinary System**

The urinary bladder is moderately distended with anechoic urine, and no luminal sediment is present. The ureteral papillae, trigone and pelvic urethra (visible to 3.0 cm) are of normal appearance, and the ureters are not visible (normal). No masses, calculi or mucosal irregularities are noted.

**SEX**

Spayed Female

The kidneys are of normal size and shape and exhibit appropriate corticomedullary differentiation with a normal 1:3 cortex to medulla ratio. There is no evidence of nephrolithiasis, mineralization, pyelectasia, cystic change or hydronephrosis. The proximal ureter is not visible (normal). The left kidney measures 6.7 cm. The right kidney measures 6.4 cm.

**AGE**

2 Years

**Adrenal Glands**

The adrenal glands are both identified in their normal locations. They are normal in size and shape with appropriate parenchymal echogenicity and normal phrenic vasculature. The left adrenal gland measures 5.7 mm at the cranial pole and 6.3 mm at the caudal pole. The right adrenal gland measures 1.5 mm at the cranial pole and 7.4 mm at the caudal pole.

**WEIGHT**

30 kg

**Spleen**

The spleen is of appropriate size and has a normal, homogenous parenchyma with a smooth, continuous capsular surface. The splenic vasculature is normal with no evidence of congestion or thrombosis, and blood flow through the splenic hilus appears normal.

**INTERPRETED BY**

Tam Mengine, DVM,  
DABVP (canine/feline  
practice)

**Liver**

The liver is of appropriate size and shape, with sharp borders and a mildly coarse parenchymal echotexture that is hypoechoic to the spleen. The portal and hepatic vasculature are of normal size and appearance with no evidence of congestion or thrombosis.

**IMAGING PERFORMED BY**

Kelly Reschny

**HOSPITAL NAME**

BPH Ancaster

The gallbladder is moderately distended with anechoic contents. The wall was thin and continuous with no focal lesions. The cystic and common bile ducts are normal / not visible.

**Gastrointestinal**

**REFERRING VET**

Dr. Pandya

The stomach is empty. The gastric wall is normal in thickness (4.6 mm) with normal deviations due to rugal folds, and exhibits appropriate wall layering. The pylorus is of normal appearance.

**INVOICE**


40824

The visualized portions of the duodenum, jejunum, and ileum are of normal thickness with intact wall layering that exhibits the appropriate 1:3 muscularis to mucosa ratio. Jejunum wall measures up to 4.2 mm. Duodenum wall measures up to 5.3 mm. Intestinal motility appears normal.

**DATE**

8/29/22

The visible portions of the colon are of normal thickness (1.6 mm) with intact wall layering. The ileocecal junction is seen and is normal.



**PATIENT** *Pancreas*

Savy Busch The areas of the limbs and body of the pancreas are isoechoic to the surrounding mesenteric fat, with normal capsular appearance. There is no evidence of peripancreatic inflammation. The pancreatic duct appears normal.

**SPECIES** *Free Abdomen*

Canine There is no evidence of free fluid within the peritoneal cavity. The omentum and intra-abdominal fat are of appropriate echogenicity. Enlarged abdominal lymph nodes are not observed. The aortic trifurcation has normal blood flow with no evidence of thrombosis.

**BREED** *ULTRASONOGRAPHIC FINDINGS*

Weimaraner X  

- Unremarkable canine abdomen

**SEX** *INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS*

Spayed Female There is no apparent cause for the ongoing gastrointestinal signs on today's ultrasound. Although the adrenal glands appear normal, screening for Addison's disease is a consideration. Additional recommendations include:

- AGE** 2 Years
- ❖ fecal parasite testing and empiric fenbendazole treatment
  - ❖ probiotic therapy
  - ❖ bland diet
  - ❖ treatment with parenteral fluids, antiemetics, antacids and gastroprotectants as clinically indicated.
  - ❖ If signs persist, trials with a novel protein or hydrolyzed diet, a resting cortisol level and a GI panel could be considered.

**INTERPRETED BY**

Tam Mengine, DVM,  
DABVP (canine/feline  
practice)

**IMAGING PERFORMED BY**

Kelly Reschny

**HOSPITAL NAME**

BPH Ancaster

**REFERRING VET**


Dr. Pandya


**INVOICE**

40824

**DATE**

8/29/22





**PATIENT**

Savy Busch

**SPECIES**

Canine

**BREED**

Weimaraner X

**SEX**

Spayed Female

**AGE**

2 Years

**WEIGHT**

30 kg

**INTERPRETED BY**

Tam Mengine, DVM,  
DABVP (canine/feline  
practice)

**IMAGING  
PERFORMED BY**

Kelly Reschny

**HOSPITAL NAME**

BPH Ancaster

**REFERRING VET**


Dr. Pandya

**INVOICE**

40824

**DATE**

8/29/22



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Tam Mengine, DVM, DABVP (canine/feline practice)

info@SonoPath.com